

**Research paper**

# **Leaf macronutrient composition of mulberry (*Morus indica*) varieties and its relationship with productivity of mulberry silkworm (*Bombyx mori* L.)**

**Kedir Shifa<sup>\*</sup>, Metasebia Terefe, Abiy Tilahun, Ahmed Ibrahim and Yohannis Nigusu**

Melkassa Agricultural Research Center, P.O.Box 436, Adama, Ethiopia

*\*Corresponding Author: Kedir Shifa, Melkassa Agricultural Research Center, P.O.Box 436, Adama, Ethiopia, Tel: +251911384487, E-mail: kedirsh01@gmail.com*

Accepted 27 May 2020

The growth, yield and reproduction potential of silkworms can be affected by their feed sources to a great extent. Minerals are among important biochemical components of leaves and they may have very big influence in productivity of silkworms. In the present study, six different mulberry varieties (K-2, S-13, M-4, Nekemte coll, Jimma Coll and Local) were evaluated for their leaf macro-mineral composition and effect on mulberry silkworms (*Bombyx mori* L.) in Melkassa Agricultural Research Center (MARC), East- Shewa zone of Oromia Regional State, Ethiopia. Completely Randomized Design (CRD) with three replications was employed for carrying out feeding trial in a laboratory. The leaf macro-mineral composition of mulberry varieties for nitrogen, phosphorus, potassium, calcium, magnesium and sulphur was determined by using appropriate methodologies. The relationship of mineral components of leaves with silkworm parameters was also determined by using correlation coefficient analysis. Overall data analysis was carried out by using SAS software at 5% probability level. Finally, the leaves of these varieties showed significant differences among themselves in their mineral compositions. Their effect on rearing performance of mulberry silkworms was also statistically different when they were fed. So, relationship of macro-mineral constituents of mulberry varieties with rearing performance of mulberry silkworms was worked out. Hence, positive correlation of nitrogen, phosphorus, potassium, calcium, magnesium and sulphur was obtained with larval, cocoon and grainage parameters of mulberry silkworms. Therefore, it can be concluded that presence of these macro-minerals in mulberry leaves will determine to the gains on important mulberry silkworm parameters when these leaves served as feed sources. As a result, these macro nutrients will be considered as basic parameters for evaluation of mulberry varieties for mulberry silkworms rearing in the future.

**Key words:** Mulberry silkworms, *Bombyx mori*, mineral composition, correlation, performance

**Cite this article as:** Kedir S, Metasebia T., Abiy T., Ahmed I., Yohannis N (2020). Leaf macronutrient composition of mulberry (*Morus indica*) varieties and its relationship with productivity of mulberry silkworm (*Bombyx mori* L.). Acad. Res. J. Agri. Sci. Res. 8(4): 390-398

## INTRODUCTION

Silk is a functional term used to describe protein fibers that are produced by arthropods. It is a natural protein fiber and is very soft, lustrous, smooth, strong and durable than any natural or artificial fibre (Lamelu, 1998; Hiwar, 2001 and Craig and Riekel, 2002). The industrial use of silk and its economic importance of production finely contributed to the silkworm promotion all over the world (Ramesh-Babu et al., 2010).

Tzenov (2007) indicated that there is a good potential for sericulture development, not only in the East Asia, but also in Eastern Europe, Central Asia, Latin America and the Africa Region based up on overall socio-economic and agro-climatic conditions (rainfall, soil, temperature, humidity, light and air). Hence, the developing countries are directing their development strategies on the productivity levels in rural areas (Hajare *et al.*, 2007). In Ethiopia, silk has played an important part in the social and religious life of Ethiopia from the earliest days of its history. This silk was imported in large quantities from India, Arabia and China and stored in vast caverns in the central highlands of Ethiopia and Ethiopian Emperors would make prodigious gifts of silk to recognized organizations and special guests. Ceremonial umbrellas, binding of sacred books, covers for wooden altars and spectacular hangings have all been produced from silk over the centuries (Spring and Hudson, 2002). Recently, silk production from mulberry silkworm (*S. c. ricini*) is practiced in Ethiopia (Metaferia *et al.*, 2007).

The mulberry-silkworm, *Bombyx mori* L., is the most exploited and commercialized silkworm. It is a sericigenous insect exploited for its valuable mulberry silk. It could be classified as multivoltine, bivoltine or univoltine based on generations per year. However, it is a monophagus species feeding only on mulberry plant and it can be reared throughout the year depending on the availability this feed plant. The agro-ecology and feed availability are the major requirements which have significant effect on rearing of larvae of this insect and finally cocoon crop yield and quality (Debaraj et al., 2003 and Thangavelu and Phukon, 1983).

Mulberry (*Morus spp.*) can ensure production of good quality cocoons. However, mulberry shows a wide range of variability in nature. It has been well recognized that morphological features and nutritive values of the leaves differ significantly from variety to variety and over locations. The morphological traits and nutritional factors could be used in selection of varieties (Solanki and Joshi, 2001). In addition, the quality of feeds given to insects generally affects the economic products of insects. So, silkworm nutrition is one and the major factor which affect development and productivity of silkworms (Milner, 2004 and Baruah and Baruah, 2007). Silkworm nutrition deals about the substances required by silkworm for its growth and metabolic functions and obtained from ingested food

and some other nutritional components synthesized through various biochemical pathways including proteinous silk fiber of commercial interest (Takano and Arai, 1978 and Hamano *et al.*, 1986).

Silkworms have adapted to feeding to derive and store adequate energy, nutrients and water from the food they consume. Feeding silkworms provides energy for growth, development, reproduction and many of its other needs. Such nutritional requirements in food consumption have direct impact on the overall genetic traits such as larval and cocoon weight, amount of silk production, pupation and reproductive traits. Thus, successful sericulture depends on increased production of leaves with high nutritive values. Therefore, the better the quality of leaves, the greater the chances of getting good cocoon harvest (Krishnaswami, 1978).

Moreover, a significant correlation has been reported between chemical compositions of mulberry leaf and larval growth and cocoon characters. Nutritional value of food plants either alone or in combination plays an important role in the larval growth and silk productivity. Improvement in larval and cocoon characters has also been witnessed with increase in chemical composition of leaves (Sannappa and Jayaramaiah, 1999).

Therefore, feed stuffs can be analyzed in to different fractions including the ash (minerals). The ash, residue left behind after all organic matter has been burnt off, is a measure of the total content of mineral salts in a food. Minerals including macronutrients constitute very good fraction of silkworm feeds. The role of inorganic salts (minerals) is variable. It has been reported that the silkworm larval structure at different ages utilizes these absorbed minerals (Ito, 1978 and Friend, 1958). Their role in the egg stage is also recognized to regulate the hydrogen ion concentration osmotic pressure. Furthermore, sodium and potassium ions act on protein synthesis. However, the importance of nitrogen has been significantly considered in different stages together with the protein content (Legay, 1958). Among minerals, nitrogen, phosphorus, potassium, magnesium, calcium, iron, silicon, sulphur, aluminium, manganese and copper found orderly on the quantity basis from different feeds of silkworms (Fukuda *et al.* 1960; Maynard and Loosli, 1962 and Sannappa and Jayaramaiah, 1999). In another way, Jayaramaiah and Sannappa (1998) find out a positive significant relationship between foliar constituents and important silkworm components. They indicated that larval duration, weight, survivability and effective rate of rearing (ERR) as well as cocoon yield were found to have significant positive relationship with nitrogen, phosphorus, potassium, calcium, magnesium and sulphur.

Therefore, selection of mulberry varieties by considering macronutrients is an important criterion for better growth and development of mulberry-silkworm (Joshi and Misra, 1982). However, very little information is available regarding such cases in Ethiopia. This work,

therefore, attempts to assess the extent of macro-mineral composition of some mulberry varieties and find out the relationship with productivity of mulberry silkworm, *Bombyx mori*.

## MATERIALS AND METHODS

### Description of the study area

The study was conducted in Melkassa Agricultural Research Center (MARC). It is found 117 kms away from Addis Ababa and 17 km to southeast of Adama in the East-Shewa zone of Oromia region, Ethiopia. It is located 8°24'N latitude and 39°12'E longitude having an elevation of 1500 meters above sea level and a mean annual rainfall of 770 mm.

### Foliar nutritional evaluation of mulberry varieties

Six selected varieties of mulberry viz. K-2, S-13, M-4, Nekemte coll, Jimma Coll and Local (check) were used in the study. These materials were selected from preliminary research observations. The methods of Sannappa and Jayaramaiah (1999) were adopted for foliar nutritional analysis on the above mentioned mulberry varieties. The leaf samples were collected after fifteen months of planting at three different heights of the plant viz. top, middle and bottom leaves in paper bags. They were shade dried and then transferred to hot air oven maintained at 70°C until constant weight was obtained. Then, the dried-up leaf samples were grinded in to fine powder and preserved for chemical analysis.

By adopting the techniques and procedures, analysis of mineral compositions for nitrogen, phosphorus, potassium, calcium, magnesium and sulphur contents were carried out for each mulberry variety. To get the estimates, dry ashing (Total Minerals) and Sample preparation were done at Melkassa Food science Laboratory, Melkassa, Ethiopia. Minerals quantification reading using ICP-OES were done at Horticoop Ethiopia (Horticultural) Laboratory, Bishoftu, Ethiopia. During analysis of aqueous solutions by ICP-OES with radial plasma observation, all measurements were performed

with spectro arcs optical emission spectrometer optimized with small volume and 32 linear CCD detectors the wave length range between 130-770 nm simultaneously analyze, the sample is nebulizer in to the argon filed plasma and the energy excitement is processed in the single spectra and the energy is expressed in intensity which is directly proportion to concentration, the concentration is calculated on the linear graph of the standard concentration and the corresponding intensities (Haynes, 1980).

### Evaluation of mulberry varieties on rearing performance of mulberry silkworms

The above mentioned mulberry varieties were used as feed source of mulberry silkworms in a laboratory. As per the rearing recommendations of silkworms by Dayashankar (1982), a bivoltine mulberry-silkworm breed was used for this experiment and it was reared on the six mulberry varieties. The silkworm rearing room and equipments were cleaned, washed and disinfected with 2% formalin solution at the rate of 800ml per 10m<sup>2</sup> before the commencement of the experiment (rearing). This breed was reared following cellular rearing techniques starting from brushing till cocoon spinning on the six varieties. Tender leaves of mulberry were fed four times a day until the larvae ends II instar stage, and semi tender leaves to III instar while more matured leaves were fed to IV and V instar larvae. The experiment was arranged in Completely Randomized Design (CRD) in three replications. In each replication, 100 worms were used and allowed to complete the larval period. As used by Neupane *et al.* (1990), Singh and Benchamin (2002) and Ramesha *et al.* (2010) matured worms were picked and mounted on the mountages for spinning. On the six day of spinning, the cocoons were harvested, counted and weighed.

Rearing parameters like larval duration (in hours), body weight of mature larvae (g), percentage of larval mortality and hatchability, cocoon traits (cocoon weight, shell weight and silk ratio), effective rate of rearing (%) and fecundity (number of eggs per female) were recorded to find out rearing performance of the worms on the mulberry varieties. As a result the following formulae were adopted for estimation when appropriate.

The following formulae, which were adopted by Singh and Benchamin (2002) used for data on rearing performance:-

$$\text{Hatching percent} = \frac{(\text{Number of normal eggs} - \text{number of non hatched eggs}) * 100}{\text{Number of normal eggs}}$$

$$\text{Effective rate of rearing (ERR)} = \frac{(\text{No. of larvae spinning cocoon} * 100)}{\text{No. of larvae brushed}}$$

## Data Analysis

Finally, the data for significance determination were analyzed statistically (one-way CRD) by using SAS software at 5% level of significance. Moreover, correlation coefficients of minerals of leaves of the mulberry varieties with their corresponding rearing performance and cocoon traits of silkworms was carried out to know their inter-relationship similarly by SAS software at 5% level of significance (SAS Institute Inc,1999-2000) .

## RESULTS

Substantial variability has been obtained in mineral composition of mulberry varieties and these differences also caused to variability in rearing performance of mulberry silkworms when served as feed sources.

### Mineral composition of different mulberry varieties

All mulberry varieties differed significantly from one another in respect to minerals (nitrogen, phosphorus, potassium, calcium, magnesium and sulphur) considered in this study. Mean values of foliar compositions of the varieties are presented in table 1.

#### A. Nitrogen

The nitrogen content of the leaf material of mulberry varieties was significantly different. It was highest in varieties of K2 (2.02582 %) followed by S13 (1.9021 %) but lowest in local (1.42232%).

#### B. Phosphorus

Phosphorus concentration of leaves of mulberry varieties was significantly different from each other with minimum record of 1107.51 mg/kg Jimma coll and maximum record of 3223.70 mg/Kg in M-4.

#### C. Potassium

The potassium content of mulberry varieties ranges from 11349.28 mg/kg in local check to 18610.38 mg/kg in M-4.

#### D. Calcium

Calcium concentration of mulberry varieties has also shown significant difference and ranges from 13452.40 mg/kg (local check) to 20522.80 mg/kg (M-4).

#### E. Magnesium

Magnesium concentration of mulberry varieties was minimum in local check (871.60 mg/kg) and maximum in M-4 (1948.93 mg/kg) and S-13 (1735.903 mg/kg).

#### F. Sulfur

The sulfur concentration of mulberry varieties was also worked out and it ranged from 152.89 mg/kg in K-2 to 338.337 mg/kg in M-4. The second highest sulfur concentration was 288.537 mg/kg in S-13.

**Table 1:** - Macronutrient (mineral) composition of different mulberry varieties

Mulberry varieties	N (mg/kg)	P (mg/kg)	K (mg/kg)	Ca (mg/kg)	Mg (mg/kg)	S (mg/kg)
Nekemte coll	1.66600c	1157.45e	12578.59d	14293.1d	1393.597c	169.170e
Jimma coll	1.70800c	1107.51f	13332.55c	15196.2c	1312.130d	187.420d
M-4	1.72200c	3223.70a	18610.38a	20522.8a	1948.930a	338.337a
K-2	2.02582a	1333.31d	13466.79c	16212.7b	1276.403e	152.890f

S-13	1.90210b	2982.39b	16030.17b	15513.7c	1735.903b	288.537b
Local	1.42232d	1731.03c	11349.28e	13452.4e	871.600f	246.240c
SE	0.04665276	208.896348	584.952079	550.245407	83.6498432	16.2461907
CV (%)	2.320974	0.918436	0.855966	1.425278	0.804390	1.310189
Pr	<.0001	<.0001	<.0001	<.0001	<.0001	<.0001

Note: - Means with the same letter are not significantly different at Pr > 0.05.

### Effect of different mulberry varieties on rearing performance

In this experiment, all the characters of mulberry silkworm rearing like hatching percentage, larval duration, larval weight, single cocoon weight, single shell weight, shell (silk) ratio, larval duration, fecundity, effective rate of rearing (ERR) and survival and showed significant differences when mulberry silkworm fed with different mulberry varieties. The results of silkworm rearing characters fed on different mulberry accessions are summarized in the table 2.

#### A. Fecundity

Silkworm fed on S-13 showed significantly higher fecundity (305.467) followed by M-4 (289.867). The local check (259.533) was lower performing compared to the rest mulberry varieties for fecundity of mulberry silkworms.

#### B. Percentage hatching of eggs

Hatching rate of silkworm eggs in to larva showed significant variation when a silkworm fed on different mulberry varieties ranging from 62.02 % to 75.33 %. The maximum hatching was recorded in M-4 (75.33 %) followed by K-2 (70.7033 %) and S-13 (70.00 %) (see Table 2).

#### C. Larval weight

Matured worms fed on S-13 recorded significantly higher larval weight (2.5303 g) and 8.17g respectively. However, the least larval weight was obtained from worms fed on local check (2.00633 g).

#### D. Effective rate of rearing (ERR)

ERR has also revealed a significant difference when mulberry silkworm fed on different mulberry varieties. Mulberry silkworm fed on S-13 (81.50 %) recorded higher ERR closely followed by M-4 (77.0 %) and Jimma (74.02 %). The least ERR was obtained from local check (68.833 %) (see Table 2).

#### E. Cocoon weight and shell weight

In accordance to yield, maximum single cocoon weight was recorded in those larvae which were fed on the leaves of S-13 (1.11167 g). The minimum single cocoon weight (0.86567 g) was recorded from larvae fed on local check. Similarly, single shell weight of a cocoon revealed significant variation when fed with the different mulberry accession. S-13 revealed significantly higher shell weight (0.187 g) but local check showed the least shell weight (0.136 g).

**Table 2:** - Effects of mulberry varieties on performance of a bivoltine mulberry silkworms

Treatment	Fecundity (%)	Hatching percent	Larval weight (gram)	ERR (%)	Cocoon weight (gram)	Shell weight (gram)
K2	279.933c	70.7033b	2.17700b	75.56667c	1.03000b	0.16833ab
S13	305.467a	70.0000b	2.53033a	81.50000a	1.11167a	0.18700a
M4	289.867b	75.3300a	2.17100b	77.00000b	1.01333b	0.16333ab

Nekemte	270.533c	65.4300c	2.19200b	73.67000e	1.00133	0.15867bc
Jimma	278.200c	62.0200c	2.23300b	74.02000d	1.00600b	0.16133bc
Local	259.533d	65.0000c	2.00633c	68.83333f	0.86567c	0.13600c
SE	3.67579869	1.07154353	0.04213927	0.92527588	0.01868696	0.00461825
CV (%)	1.973116	0.637592	4.154814	0.157055	3.194023	8.825915
Pr	<.0001	<.0001	0.0005	<.0001	<.0001	0.0238

Note: - Means followed by the same letter within a column are not significantly different from each other at 5% level of probability (LSD mean comparison test)

### Relationship of mineral constituents of leaves of mulberry varieties with mulberry silkworm rearing performance

The foliar mineral composition of mulberry varieties resulted in significantly important inter-relationship with performance of mulberry silkworms when their leaves served as feeds. This relationship between leaf composition values and important mulberry silkworm traits have been worked out through correlation analysis and the results are summarized in table 3.

#### Nitrogen

It revealed significantly strong positive correlation with larval weight ( $r= 0.56058$ ), effective rate of rearing ( $r= 0.74339$ ), cocoon weight ( $r=0.77688$ ) and shell weight ( $r=0.64814$ ). However, its relationship (correlation) was positive to all silkworm characters though not statistically significant.

#### Phosphorus

Phosphorus content of leaves had shown significant positive correlation with egg fecundity ( $r=0.66843$ ) and hatchability ( $r=0.76540$ ) as well as larval effective rate of rearing ( $r=0.61648$ ). However, it had shown no significant correlation with cocoon characters of cocoon and shell weight.

#### Potassium and Calcium

Potassium composition of mulberry leaves revealed significantly positive correlation with egg fecundity ( $r=0.75357$ ), hatchability ( $r=0.81335$ ), larval effective rate of rearing ( $r=0.74040$ ) and cocoon weight ( $r=0.55303$ ). Moreover, calcium was with significant positive correlation with w with egg fecundity ( $r=0.50507$ ) and hatchability ( $r=0.82553$ ) as well as larval effective rate of rearing ( $r=0.47879$ ).

#### Magnesium and sulphur

Magnesium have been found to have significant positive correlation with all mulberry silkworm rearing characteristics under study at  $p < 0.05$ , which include egg fecundity, hatchability, larval weight, effective rate of rearing, cocoon weight and shell weight. However, sulphur was significantly positively correlated only with egg parameters of egg fecundity ( $r=0.47178$ ) and hatchability ( $r=0.60089$ ).

**Table 3:** - Relationship of minerals with mulberry silkworm rearing performance

Minerals	Silkworm traits					
	Fecundity	Egg hatchability	Larval weight	Effective rate of rearing	Cocoon weight	Shell weight
<b>N</b>	0.65510*	0.45498	0.56058*	0.74339*	0.77688*	0.64814*
<b>P</b>	0.66843*	0.76540*	0.38065	0.61648*	0.34554	0.33381
<b>K</b>	0.75357*	0.81335*	0.41267	0.74040*	0.55303*	0.46559
<b>Ca</b>	0.50507*	0.82553*	0.09385	0.47879*	0.33893	0.27559
<b>Mg</b>	0.79270*	0.70851*	0.54266*	0.82974*	0.69242*	0.56857*
<b>S</b>	0.47178*	0.60089*	0.19262	0.37572	0.10033	0.11116

\* Significant at  $Pr \leq 0.05$

## DISCUSSION AND CONCLUSION

It can be realized from the results of the present investigation that mulberry varieties (K-2, S-13, M-4, Nekemte coll, Jimma Coll and local check) showed wide variation in their qualitative traits with respect to minerals from such studies of the first type in Ethiopia. Consequently, since accumulation of nutrients in insect greatly influenced by the nutritional richness of the host plants, the macro-mineral compositions (nitrogen, phosphorus, potassium, calcium, magnesium and sulphur) of leaves of mulberry varieties have been studied and they revealed very significant differences among mulberry varieties.

Similarly, studies on different mulberry varieties have been conducted in abroad by several authors and they have found out significant differences among different varieties (Jayaramaiah and Sannappa, 1998). In addition, (Scriber and Slansky, 1981) emphasized the importance of nitrogen for silkworm rearing. Accordingly, nitrogen was comparatively higher in genotype K-2 and S- 13 and from this study, which recorded 2.026% and 1.902%, respectively. The average nitrogen content obtained from this study was lower than the findings of Murthy et al (2013) which could be due to the soil conditions and the delayed timing of nutrient analysis.

Furthermore, the inter-relationship of performance of mulberry silkworms and the macronutrient composition of leaves of mulberry varieties have been analyzed. The results suggest that different traits of mulberry silkworm have significantly different relationship with various biochemical constituents of mulberry leaves. As a result, cocoon traits such as cocoon weight and shell weight found to have significant positive relationship with nitrogen, potassium and magnesium. In addition, larval characteristics like larval weight and effective rate of rearing showed a significant positive correlation or interaction with nitrogen, phosphorus, potassium, calcium and magnesium. Moreover, grainage parameters like egg fecundity and hatching showed significant positive

correlation with phosphorus, potassium, calcium, magnesium and sulphur composition of mulberry leaves. In general, the significant positive relationship obtained from foliar macronutrient compositions (nitrogen, phosphorus, potassium, calcium, magnesium and sulphur contents) and silkworm performance indicators was found to be very important for consideration of mulberry varieties as obtained Jayaramaiah and Sannappa (1998) and Murthy *et al* (2013).

Hence, the present study reveals that the selection of mulberry varieties for rearing mulberry silkworm based up on foliar macro-mineral constituents of the mulberry varieties as well as rearing performance is very important in order to get improvements in larval development, cocoon production and grainage parameters. However, more research should also be carried out to support the current findings in consideration with varying period of leaf picking and nutrient analysis, leaf proximate composition, field performances of mulberry varieties and in relation feeding efficiency of mulberry silkworms when using these varieties as feed sources.

## REFERENCES

- A.O.A.C. (1970). *Methods of Analysis*, 9<sup>th</sup> edn. Association of Official Agricultural Chemists, Washington, 789pp.
- Baruah, A.B. and Baruah, P.K. (2007). Studies on the status of biochemical constituents in four morphotypes of muga food plant som (*Persea bombycina*). In: *Proceedings of International Conference on "Sericulture Challenges in the 21<sup>st</sup> Century and the 3rd BACSA meeting*, September 2007, Vratza.
- Craig, C. L. and Riekel, C. (2002). Comparative architecture of silks, fibrous proteins and their encoding genes in insects and spiders. *Comparative Biochemistry and Physiology*. **133**: 493-507.
- Dayashankar, K.N. (1982). Performance of eri silkworm, *Samia cynthia ricini* Boisduval on different

- host plants and economics of rearing on castor under Dharwad conditions. *M.Sc. Thesis*, University of Agricultural Sciences, Bangalore, pp.60-86.
- Debaraj, Y., Singh, B.K., Das, P.K. and Suryanarayan, N. (2003). Payam: An evergreen host plant of eri silkworm. *Indian silk* **5**:5-6.
  - Devaiah, M.C., Rajashekhargouda, R., Suhas, Y. and Govindan, R. (1985). Growth and silk production in *Samia cynthia ricini* Boisduval as fed on four different host plants. *Ind.J. Seric* **24**: 33-35
  - Friend, W.G. (1958). Nutritional requirements of phytophagous insects. *Ann. Rev. Ent.* **3**: 57-74.
  - Fukuda, T., Kamegame, T. and Matsuda, M. (1963). A correlation between the mulberry leaves consumed by the mulberry silkworm larvae in different ages of the larval growth and production of cocoon fiber spun by silkworm larvae and the eggs laid by the silkworm. *Bull. Seric. Exp. Stn.* **8**: 165-171.
  - Fukuda, T., Suto, M. and Higuchin, Y. (1960). Silkworm raising on the artificial food. *J. Seric. Sci.* **29**:1-13.
  - Hajare, T.N., Jadhav, A.D, Venugopalan, M., Patil, N.G., Chaturvedi, A. and Maji, A.K. (2007). Evaluation of sericulture for augmenting agricultural income of marginal farmers in semi-arid region of India. In: *Proceedings of International Conference on Sericulture Challenges in the 21<sup>st</sup> Century and the 3rd BACSA meeting*, September 2007, Vratza.
  - Hajarika, U., Barah, A., Phukan, J.C. and Benshamin, K.V. (2003). Study on the effect of different food plants and seasons on the larval development and cocoon characters of silkworm *Samia cynthia ricini* Boisduval. *Bull. Ind. Acad. Seric.* **7**:77- 85.
  - Hamano, K., Miyazawa, K. and Mukiyama, F. (1986). Racial difference in the feeding habit of the silkworm, *Bombyx mori*. *J. Seric. Sci.* **55**: 68-72.
  - Hiwar, C.J. (2001). *Agro Cottage Industry: Sericulture*. Daya Publishing House. New Delhi. pp. 1-117.
  - Haynes, R. J. (1980). A Comparison of Two Modified Kjeldahl Digestion Techniques for Multi-Element Plant Analysis with Conventional Wet and Dry Ashing Methods. *Comm. in Soil Sci. and Plant Anal.* **11**(5): 459-467.
  - Ito, T. (1978). Silkworm nutrition. In: *The Silkworm: an Important Laboratory Tool*, pp. 150–151 (Tazima, Y., ed). Kodansha Ltd., Tokyo.
  - Jayaramaiah, M. and Sannappa, B. (1998). Correlation Coefficients Between Foliar Constituents of Castor Genotypes and Economic Parameters of the Eri Silkworm, *Samia cynthia ricini* Boisduval (Lepidoptera: Saturniidae). *The third Int. Conf. Wild Silkmooths*, pp. 162-194.
  - Joshi, K.L. and Misra, S.D. (1982). Silk Percentage and effective rate of rearing of eri silkworm, *Philosamia ricini* Hutt. (Lepidoptera: Saturniidae). *Entomol.* **7**:107-110
  - Krishnaswami, S. (1978). New Technology of Silkworm Rearing. *Bull.Cent. Seric. Res.Train. Inst., Mysore*, pp. 1-24.
  - Lamelu, A.G. (1998). *Illustrated textbook on sericulture-translated from Japanese*. Siences pulishing Inc. Enfield, New Hampshire, pp.10-35
  - Legay, J. M. (1958). Recent advances in silkworm nutrition. *A. Rev. Ent.*, **3**: 75-86.
  - Maynard, A.L. and Loosli, L., (1962). *Animal Nutrition*, 5<sup>th</sup> edn.. McGraw Hill Book Co., Inc., New York, pp: 533.
  - Metaferia, H.Y., Amanuel, T. and Kedir, S. (2007). Scaling up of silk production technologies for employment and income generation in Ethiopia. In: *Success With Value Chain: proceedings of scaling up and scaling out of agricultural technologies in Ethiopia, an international conference, 9-11 May 2006* (Tsedeke Abate ed). Ethiopian Institute of Agricultural Research, Addis Ababa.
  - Milner, J.A. (2004). Nutrition and gene regulation: Molecular targets for bioactive food components. *J. Nutr.* **134**: 2492-2498.
  - Murthy, V.N.Y, Ramesh, H.L. Lokesh, G., Munirajappa and Yadav, B.R.D (2013). Leaf Quality Evaluation of Ten Mulberry (*Morus*) Germplasm Varieties through Phytochemical Analysis. *Int. J. Pharm. Sci. Rev. Res.*, **21**(1): 182-189
  - Neupane, F.P., Thapa, R.B. and Parajulee, M.N. (1990). Life and seasonal histories of the eri silkworm, *Samia cynthia ricini* Hutt. (Lepidoptera: Saturniidae) in Nepal. *J. Inst. Agri.: Anim. Sci.* **11**:113–120.
  - Patil, G.M., Kulkarni, K.A., Patil, R.K. and Badiger, K.S. (1998). Performance of Eri Silkworm, *Samia Cynthia Ricini* Boisd. on different castor genotypes. *Int. Conf. Wild Silkmooths*. PP. 193-195
  - Patil, R.R., Kusugal, S. and Ankad, G. (2009). Performance of eri silkworm, *Samia cynthia ricini* Boisd. on few food plants. *Karnataka J. Agric. Sci.* **22**: 220-221.
  - Ramesha, C., Anuradha, C.M., Lakshmi, H., Kumari, S.S., Seshagiri, S.V. Goel, A.K. and Kumar, C.S. (2010). Nutrigenetic traits analysis for the identification of nutritionally efficient silkworm germplasm breeds. *Biotechnology* **9**: 131-140.
  - Sahlemedhin Sertsu and Taye Bekele (2000). *Procedures for soil and plant analysis*. Technical paper, Ethiopian Agricultural Research Organization, Addis Ababa, pp. 60-83
  - Sannappa, B. and Jayaramaiah, J. (1999). Mineral Constituents of Selected Genotypes of Castor, *Ricinus communis* L. *Mysore J.agric.Sci.* **33**: 157-161
  - SAS Institute Inc (1999-2000). *SAS/STAT users' guide*. SAS Institute. Cary.
  - Scriber, J. M. and Slansky, F. (1981). The Nutritional Ecology of Immature Insects. *Ann. Rev. Entomol.* **26**: 183-211
  - Singh, K.C. and Benchamin, K.V. (2002). Biology and ecology of the eri silkworm *Samia ricini* Donovan



- (Saturniidae): *Bullet. Ind. Acad. Seric.* **6**: 20-33.
- Solanki, S.S. and Joshi, P. (2001). Inheritance study of some morphological traits in castor (*Ricinus communis* L.). *Ind. J. Genet.* **61**:136-139.
  - Spring, C. and Hudson, J. (2002). *Silk in Africa: Ethiopia*. University of Washington Press, Seattle.
  - Thangavelu, K. and Phukon, J.C. (1983). Food Preference of Eri Silkworm *Philosamia ricini* Hutt (Saturnidae: Lepidoptera). *Entomol.* **8**: 311-315
  - Tzenov, P. (2007). Present status and utilization of sericulture germplasm and comparative studies of different silkworm hybrids performance for sericultural enterprise development in the Black, Caspian seas and Central Asia (BACSA) region. **In**: *Proceedings of International Conference on Sericulture Challenges in the 21<sup>st</sup> Century and the 3rd BACSA meeting, September 2007, Vratza*